

Spectroscopic studies of the vibrational and electronic properties of hydrogen at high pressure- temperature conditions

*Alexander F. Goncharov, Eugene Gregoryanz,
Russell J. Hemley, Ho-kwang Mao*

This article was submitted to International Symposium on
Advanced Materials, Tsukuba, Japan,
March 10-13, 2003

U.S. Department of Energy

Lawrence
Livermore
National
Laboratory

DISCLAIMER

This document was prepared as an account of work sponsored by an agency of the United States Government. Neither the United States Government nor the University of California nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or the University of California, and shall not be used for advertising or product endorsement purposes.

This is a preprint of a paper intended for publication in a journal or proceedings. Since changes may be made before publication, this preprint is made available with the understanding that it will not be cited or reproduced without the permission of the author.

This work was performed under the auspices of the United States Department of Energy by the University of California, Lawrence Livermore National Laboratory under contract No. W-7405-Eng-48.

This report has been reproduced directly from the best available copy.

Available electronically at <http://www.doc.gov/bridge>

Available for a processing fee to U.S. Department of Energy
And its contractors in paper from
U.S. Department of Energy
Office of Scientific and Technical Information
P.O. Box 62
Oak Ridge, TN 37831-0062
Telephone: (865) 576-8401
Facsimile: (865) 576-5728
E-mail: reports@adonis.osti.gov

Available for the sale to the public from
U.S. Department of Commerce
National Technical Information Service
5285 Port Royal Road
Springfield, VA 22161
Telephone: (800) 553-6847
Facsimile: (703) 605-6900
E-mail: orders@ntis.fedworld.gov
Online ordering: <http://www.ntis.gov/ordering.htm>

OR

Lawrence Livermore National Laboratory
Technical Information Department's Digital Library
<http://www.llnl.gov/tid/Library.html>

The behavior of hydrogen at high densities has been widely explored in recent years both experimentally and theoretically, yielding a wealth of information on the material (*e.g.*, Ref. 1). Detailed information has been obtained from static compression experiments generally limited to low-temperature studies (<300 K) and maximum pressures of ≈ 300 GPa (*e.g.*, Refs. [2,3]). However, there are now numerous questions regarding the behavior of hydrogen at high pressures and temperatures, results that have important implications for both fundamental physics and planetary science.

We report here [2] Raman scattering and visible to near-infrared absorption spectra of solid hydrogen under static pressure up to 285 GPa at 85-140 K. We obtain pressure dependences of vibron and phonon modes in agreement with previously determined to lower pressures. The results indicate the stability of the ordered molecular phase III to the highest pressure reached and provide constraints on the insulator-to-metal transition pressure. Extrapolations of the vibron and phonon frequencies suggest transformation to a monoatomic state below 495 GPa. On the other hand, considerations of the absorption edge indicate the pressure of metallization at 325-385 GPa on the basis of tentative extrapolation of the direct band gap energy. Although complicated by effects of stressed-induced diamond absorption and possible differences between the behavior of the direct and indirect gap, there appears to be an emerging consistence between various experimental and theoretical results, with a predicted transition at 325-495 GPa.

We also report high P - T Raman measurements of solid and fluid hydrogen to above 1100 K and to 155 GPa [4]. These conditions, which were previously inaccessible by static compression experiments, provide new insight into the behavior of the material under extreme conditions. The data give a direct measure of the melting curve (Fig. 1) that extends previous optical investigations by up to a factor of four in pressure. The measurements indicate that the melting curve may flatten out [5]. This result is close to the melting line maximum and liquid-liquid transition predicted theoretically [7,8].

The magnitude of the vibron temperature derivative $(dv/dT)_P$ increases by a factor of ≈ 30 (Fig. 2) over the measured pressure range, indicating an increase in intrinsic anharmonicity and weakening of the molecular bond. Moreover, there is an unexpected increase in intensity of the Raman vibron that arises from the combined P - T effects on the electronic properties. Overall, the results reveal significant changes in the state of hydrogen at moderate temperatures (≈ 1000 K) and megabar pressures.




Fig. 1. The melting curve of H₂. The filled circles are from the present work, open diamonds from [5] and open triangles are from [6] and references therein. Different melting laws were fitted to 3 combined data sets: dashed line - Kechin equation; dot-dashed - Kraut-Kennedy equation; dotted - Simon-Glatzel equation. The light gray line shows the Kechin model fit from Ref. datchi. Inset: phase diagram of H₂ with low temperature phases (shown in solid lines). Arrows show various P - T paths taken in the higher pressure range.




Fig. 2. A temperature derivative of the vibron frequency versus pressure. The solid line is a guide to the eye. Insert: reduced vibron frequencies as a function of temperatures for different pressures.

References:

1. J. Kohanoff, *J. Low Temp. Phys.*, **122**, 297 (2001), and references therein.
2. A. F. Goncharov, E. Gregoryanz, R. J. Hemley, and H.-K. Mao. *PNAS* **98**: 14234 (2001).
3. P. Loubeyre, F. Occelli and R. LeToullec, *Nature* (London), **416**, 613 (2002).
4. E. Gregoryanz, A. F. Goncharov, K. Matsuishi, H.-K. Mao and R. J. Hemley, *PRL*, submitted.
5. F. Datchi, P. Loubeyre, and R. LeToullec, *Phys. Rev. B* **61**, 6535 (2000).
6. V. Diatschenko *et al.*, *Phys. Rev. B* **32**, 381 (1985).
7. O. Pfaffenzeller and D. Hohl, *J. Phys.: Condens. Matter* **9**, 11023 (1997).
8. S. Scandolo, *Proc. Nat. Acad. Sci.*, in press.